© 2026 RK Publication | Volume 2, | ISBN 978-81-181100-3-3
RK PUBLICATION
Digital Nursing Book Publisher – A Unit of RK Publications
https://book.rk-publication.com/1

A Quasi-Experimental Study to Assess the Effectiveness of Foot and Hand Massage in Reducing Pain Intensity among Post-Caesarean Mothers in Selected Maternity Hospitals at Noida, Uttar Pradesh.

Prof. SHINE JOSE

Ph.D. Scholar, Desh Bhagat University, Mandi Gobindgarh, Punjab

Introduction to Post-Caesarean Pain

Caesarean section is one of the most commonly performed surgical procedures in obstetric practice. It is an operative procedure in which the fetus is delivered through an incision made in the abdominal wall and uterus. Caesarean birth may be planned or performed as an emergency depending on maternal, fetal, or obstetric indications. Although caesarean section can be a lifesaving procedure for both mother and baby, it is major abdominal surgery and is associated with postoperative pain, reduced mobility, fatigue, and temporary limitations in maternal activities.

Post-caesarean pain is an important clinical concern during the immediate postpartum period. Surgical trauma to the skin, subcutaneous tissue, muscles, fascia, and uterus stimulates nociceptive pathways and produces varying degrees of pain. The intensity of pain may differ considerably among mothers because pain is influenced not only by tissue injury but also by psychological, emotional, cultural, and social factors. A mother who experiences severe postoperative pain may have difficulty changing position, walking, coughing, deep breathing, breastfeeding, caring for the newborn, and performing personal hygiene.

Pain is defined as an unpleasant sensory and emotional experience associated with actual or potential tissue damage. In post-caesarean mothers, pain is generally acute and results primarily from surgical tissue injury. The experience may be described as incisional, pulling, burning, aching, or stabbing. Pain may increase during movement, coughing, sneezing, laughing, standing, or changing position. The mother may also experience uterine cramping associated with uterine involution.

Effective pain management is an essential component of quality postpartum nursing care. The objective of pain management is not merely to eliminate pain but to reduce pain to a level that allows the mother to participate comfortably in essential activities. Adequate pain relief promotes early ambulation, improves sleep and rest, facilitates breastfeeding, supports maternal–newborn interaction, and contributes to a positive childbirth experience.

The postpartum period represents a significant transition in a woman's life. Following caesarean birth, the mother must recover from surgery while simultaneously adapting to the physical and emotional demands of motherhood. Pain can interfere with this transition. Therefore, nurses have a central role in identifying pain, assessing its intensity and characteristics, implementing appropriate interventions, evaluating treatment outcomes, and providing education regarding pain management.

Pain management should be individualized according to the mother's clinical condition, pain severity, preferences, previous experiences, and prescribed treatment. Contemporary nursing care emphasizes a multimodal approach in which pharmacological and non-pharmacological methods may be combined. Non-pharmacological interventions such as relaxation, positioning, breathing exercises, therapeutic touch, distraction, music, heat or cold applications when appropriate, and massage may complement pharmacological treatment.

Foot and hand massage are particularly relevant nursing interventions because they are relatively simple, inexpensive, non-invasive, and potentially suitable for postpartum settings when appropriate precautions are followed. Massage may promote relaxation and comfort and may influence the perception of pain through sensory stimulation and reduction of muscle tension.

The assessment and management of post-caesarean pain should therefore be considered an integral part of maternal nursing care rather than an isolated postoperative procedure. Nurses should assess pain systematically, administer prescribed analgesics safely, provide appropriate comfort measures, encourage mobility according to clinical status, and continuously evaluate the mother's response.

Understanding Pain after Caesarean Section

Concept of Pain

Pain is a complex phenomenon involving sensory, emotional, cognitive, behavioral, and physiological components. The experience of pain differs from one individual to another. Two mothers undergoing similar surgical procedures may report different levels of pain because pain perception is influenced by previous experiences, anxiety, expectations, coping abilities, fatigue, sleep, cultural background, and individual sensitivity.

For nursing practice, pain should be considered what the patient reports experiencing. Observable behaviors such as facial grimacing, guarding, crying, restlessness, or reluctance to move can support assessment, but absence of visible behavioral signs does not necessarily indicate absence of pain.

Post-caesarean pain generally has an acute postoperative component. The surgical incision produces tissue injury, which activates peripheral nociceptors. Signals are transmitted through peripheral nerves to the spinal cord and subsequently to higher centers of the nervous system where pain is perceived.

Types of Pain Experienced after Caesarean Section

Post-caesarean mothers may experience several types of discomfort. Incisional pain occurs at the abdominal surgical site and is commonly aggravated by movement. Visceral discomfort may arise from manipulation of abdominal and pelvic structures during surgery. Uterine contractions associated with involution may produce cramping abdominal pain.

Pain can also occur at the site of intravenous cannulation, urinary catheterization, or other procedures. Musculoskeletal discomfort may result from prolonged positioning during surgery or limited mobility after surgery.

Acute Pain

Acute pain usually has a relatively recent onset and is associated with tissue injury. Postoperative caesarean pain is predominantly acute. It generally decreases as tissue healing progresses, although the rate of recovery differs between individuals.

Incisional Pain

Incisional pain originates from the abdominal surgical wound. The pain may become more noticeable during movement, coughing, sneezing, getting out of bed, or lifting the baby. Proper positioning and support of the incision during movement may improve comfort.

Uterine Cramping

After delivery, the uterus undergoes involution and contracts to gradually return toward its pre-pregnancy size. These contractions can cause cramping discomfort. The intensity may vary between mothers and may be more noticeable during breastfeeding because physiological processes associated with breastfeeding promote uterine contraction.

Physiological Basis of Postoperative Pain

Surgical tissue injury activates nociceptors. Chemical mediators released at the site of tissue injury contribute to sensitization of peripheral nerve endings. Pain impulses are transmitted through peripheral nerves to the spinal cord and then to the brain.

The nervous system does not function independently of psychological factors. Anxiety, fear, uncertainty, sleep deprivation, and fatigue may increase the mother's perception of pain. Conversely, relaxation, reassurance, adequate information, emotional support, and appropriate comfort measures may reduce distress and improve coping.

This relationship is particularly important in nursing because pain management includes both physical and psychological dimensions.

Factors Influencing Post-Caesarean Pain

Several factors may influence postoperative pain intensity.

Maternal factors include age, previous surgical experience, pain tolerance, anxiety, fatigue, sleep deprivation, and individual coping mechanisms.

Surgical factors include type of incision, duration of surgery, tissue manipulation, postoperative wound condition, and associated procedures.

Obstetric factors may include emergency versus planned caesarean birth, previous caesarean section, complications during pregnancy, and maternal or fetal conditions.

Psychological factors include fear, anxiety, stress, expectations regarding childbirth, and concerns about the newborn.

Environmental factors include noise, inadequate rest, lack of privacy, limited family support, and unfamiliar hospital surroundings.

Pain and Maternal Behavior

Pain may alter maternal behavior. A mother experiencing significant pain may avoid movement and may require assistance with basic activities. She may hesitate to hold or breastfeed the newborn because changing position is uncomfortable.

Pain may also affect communication. Some mothers may clearly report their pain, while others may minimize symptoms because they believe postoperative pain is unavoidable or because they do not want to disturb healthcare personnel. Nurses should therefore encourage mothers to report pain without embarrassment or fear.

Assessment of Pain Intensity

Pain assessment is the first major step in effective pain management. The nurse should assess pain systematically rather than relying only on observation.

A comprehensive pain assessment should include:

- Location
- Intensity
- Quality
- Onset
- Duration
- Pattern
- Aggravating factors
- Relieving factors
- Associated symptoms
- Effect on activities
- Previous pain-management methods

Importance of Pain Assessment

Accurate pain assessment provides a baseline against which the effectiveness of interventions can be evaluated. It also assists nurses and other healthcare professionals in determining whether additional intervention is required.

Pain assessment should be performed at appropriate intervals and after interventions to determine the response. The mother should be asked to describe her pain in her own words.

Numerical Rating Scale

© 2026 RK Publication | Volume 2, | ISBN 978-81-181100-3-3
RK PUBLICATION
Digital Nursing Book Publisher – A Unit of RK Publications
https://book.rk-publication.com/1

The Numerical Rating Scale is commonly used to assess pain intensity. The mother may be asked to rate pain using a numerical range, commonly from 0 to 10, where 0 represents no pain and the highest number represents the worst pain imaginable.

Score	General Interpretation
0	No pain
1–3	Mild pain
4–6	Moderate pain
7–10	Severe pain

The numerical ranges should be interpreted within the context of the particular assessment tool and clinical setting.

Visual Analogue Scale

The Visual Analogue Scale generally consists of a line representing a continuum from no pain to the worst pain. The patient indicates the point that best represents the current pain intensity. The scale can provide a quantitative measure of subjective pain.

Verbal Rating Scale

A Verbal Rating Scale asks the mother to describe pain using categories such as no pain, mild pain, moderate pain, severe pain, or very severe pain.

This method may be useful when a mother finds numerical scales difficult to understand.

Behavioral Assessment

Behavioral indicators may provide additional information. These may include facial expressions, guarding of the abdomen, reluctance to move, crying, restlessness, changes in activity, and altered interaction.

However, behavioral observation should complement rather than replace self-report in a mother who can communicate.

Assessment of Pain During Movement

Post-caesarean pain should not be assessed only while the mother is resting. Movement-related pain can significantly affect functional recovery. Nurses should assess whether pain interferes with:

- Turning in bed
- Sitting
- Standing
- Walking
- Breastfeeding
- Holding the newborn
- Coughing
- Deep breathing
- Personal hygiene

Functional assessment is especially important because the ultimate goal of pain management is to support recovery and meaningful maternal activities.

Nursing Assessment Approach

A practical nursing assessment can follow the **PQRST** approach:

P – Provocation/Palliation:
What makes the pain worse or better?

Q – Quality:
How does the pain feel? Is it aching, burning, pulling, stabbing, or cramping?

R – Region/Radiation:
Where is the pain located? Does it spread to another area?

S – Severity:
What is the intensity of pain?

T – Timing:
When did it begin? Is it continuous or intermittent?

This structured assessment helps nurses obtain comprehensive information and document pain accurately.

Impact of Postoperative Pain on Maternal Recovery

Postoperative pain can influence several aspects of maternal recovery. Effective pain control is therefore important for physical, psychological, functional, and social recovery.

Effect on Mobility

Pain may discourage mothers from moving. Prolonged immobility can delay functional recovery and may contribute to other postoperative complications. Adequate pain control can make it easier for the mother to participate in gradual mobilization as recommended by the healthcare team.

Early and safe mobilization is an important component of postoperative nursing care. Nurses should assess the mother's condition and assist her with position changes and ambulation when appropriate.

Effect on Breastfeeding

A post-caesarean mother may experience difficulty maintaining a comfortable breastfeeding position because abdominal movement can increase pain. The mother may need assistance with positioning and support.

Nurses can teach comfortable breastfeeding positions and help the mother use pillows or other appropriate supports to minimize pressure on the surgical area.

Effect on Mother–Infant Interaction

Pain, fatigue, and limited mobility may interfere with the mother's ability to hold, position, and interact with the newborn. Adequate pain management can facilitate maternal participation in newborn care.

The nurse should encourage safe mother–infant interaction while considering the mother's comfort and postoperative condition.

Effect on Sleep and Rest

Pain can interfere with sleep and relaxation. Postoperative mothers may experience repeated interruptions related to newborn care, medication administration, vital-sign monitoring, and hospital activities. When pain is added to these challenges, adequate rest may become difficult.

Nurses should therefore consider pain management as part of promoting restorative rest.

Effect on Psychological Well-Being

Persistent or severe pain can increase distress, anxiety, irritability, and feelings of helplessness. A mother who is unable to move comfortably or care for her newborn may become frustrated.

Emotional support and clear communication are important components of nursing care. The mother should be informed about expected postoperative recovery and encouraged to report pain early.

Effect on Respiratory and Physical Activities

Pain may cause a mother to limit deep breathing or coughing because these activities may increase abdominal discomfort. Appropriate support of the incision and effective pain management can facilitate participation in necessary postoperative activities.

Effect on Hospital Recovery

Uncontrolled pain may interfere with mobility, feeding, sleep, self-care, and participation in newborn care. Comprehensive pain management can support recovery and patient satisfaction.

Pharmacological Approaches to Pain Management

Pharmacological management remains an important component of post-caesarean pain control. Analgesic therapy should be administered according to the healthcare provider's prescription, institutional protocols, and the mother's clinical condition.

The choice of analgesic depends on factors such as pain severity, previous medication exposure, allergies, contraindications, renal and hepatic function, breastfeeding considerations, and other clinical factors.

Multimodal Analgesia

Multimodal analgesia refers to using different approaches or medications that act through different mechanisms to provide effective pain relief while potentially reducing reliance on any single analgesic class.

For post-caesarean mothers, analgesic plans may include non-opioid medications and, when clinically indicated, opioid medications. Regional anesthesia techniques may also contribute to perioperative pain control.

Non-Opioid Analgesics

Non-opioid analgesics are commonly used for postoperative pain management. Examples include paracetamol/acetaminophen and non-steroidal anti-inflammatory drugs when appropriate.

Paracetamol

Paracetamol is widely used for mild-to-moderate pain and may form part of a multimodal postoperative analgesic regimen.

Nurses should administer the medication according to the prescribed dose and schedule and should be aware of maximum recommended daily doses and the presence of paracetamol in combination preparations.

Non-Steroidal Anti-Inflammatory Drugs

NSAIDs reduce pain partly by inhibiting prostaglandin synthesis. They may be useful for inflammatory and postoperative pain when not contraindicated.

Nurses should monitor for potential adverse effects and contraindications, including gastrointestinal, renal, bleeding, and hypersensitivity concerns as relevant to the specific medication.

Opioid Analgesics

Opioids may be used for moderate-to-severe postoperative pain when clinically indicated. Examples include morphine and other opioid medications.

Because opioids can cause sedation and respiratory depression, careful monitoring is necessary. Other potential adverse effects include nausea, vomiting, constipation, pruritus, and dizziness.

The nurse should assess respiratory status, level of consciousness, pain response, and other relevant parameters according to the medication and institutional protocol.

Regional Analgesia

Regional anesthesia and analgesia may contribute to postoperative pain control. Techniques may include neuraxial anesthesia and regional nerve blocks depending on clinical circumstances and available expertise.

Nurses play an important role in monitoring patients following regional techniques and identifying complications requiring prompt attention.

Nursing Responsibilities in Pharmacological Pain Management

The nurse should:

1. Assess pain before medication administration.
2. Review the medication order carefully.
3. Check allergies and relevant contraindications.
4. Administer medication using safe medication practices.
5. Monitor therapeutic response.
6. Observe for adverse effects.
7. Reassess pain after an appropriate interval.
8. Document assessment, intervention, and response.
9. Educate the mother regarding prescribed medication.
10. Report inadequate pain relief or significant adverse effects.

Pharmacological treatment should not be considered the only component of postoperative pain management. Non-pharmacological interventions can be used alongside prescribed analgesia when appropriate.

Non-Pharmacological Approaches to Pain Management

Non-pharmacological pain management refers to interventions that do not primarily involve administration of analgesic medication. These approaches can support comfort, relaxation, coping, and functional recovery.

They may be particularly valuable in postpartum nursing because many techniques are simple and can be integrated into routine care.

Importance of Non-Pharmacological Methods

Non-pharmacological interventions can provide several potential benefits:

- Promote relaxation
- Reduce anxiety
- Improve comfort
- Support coping
- Encourage participation in recovery
- Complement pharmacological treatment
- Promote patient involvement in care

The intervention should be selected according to the mother's condition, preference, cultural acceptability, and clinical safety.

Positioning

Proper positioning is one of the simplest nursing interventions for reducing postoperative discomfort. The mother should be assisted to find a position that minimizes pressure on the incision and allows comfortable movement.

Pillows and appropriate supports may be used to facilitate comfortable sitting, lying, and breastfeeding.

Relaxation Techniques

Relaxation techniques can help reduce muscle tension and anxiety. Slow breathing, progressive relaxation, guided relaxation, and focused attention may be used according to the mother's ability and preference.

The nurse can provide simple instructions and encourage the mother to practice relaxation during periods of increased discomfort.

Deep Breathing

Controlled breathing may promote relaxation and help the mother cope with discomfort. The nurse can teach slow, comfortable breathing while avoiding excessive effort.

Breathing exercises may also be incorporated into routine postoperative respiratory care when appropriate.

Distraction

Distraction involves directing attention away from pain toward another activity or stimulus. Conversation, reading, music, television, relaxation recordings, or interaction with the newborn may serve as distractions depending on the mother's condition.

Music Therapy

Music may promote relaxation and reduce anxiety. The mother may select music that she finds pleasant and culturally appropriate.

Music-based interventions can be particularly easy to incorporate into patient-centered care because they require minimal equipment when personal devices are available and permitted.

Therapeutic Touch

Therapeutic touch and other forms of appropriate supportive touch may promote relaxation and comfort. Such interventions should respect the mother's consent, privacy, cultural preferences, and clinical condition.

Heat and Cold Applications

Heat or cold applications may be used for selected types of pain when clinically appropriate. However, application near a fresh surgical wound requires caution and should follow institutional guidance. The nurse should assess skin integrity, sensation, circulation, and contraindications before using these interventions.

Guided Imagery

Guided imagery involves encouraging the patient to focus attention on calming mental images. It may be used as a relaxation strategy and can complement other pain-management measures.

Foot Massage

Foot massage is a manual technique involving gentle manipulation of the feet. It may promote relaxation and comfort and can be incorporated into nursing care when there are no contraindications.

The intervention should be performed with appropriate hand hygiene, patient consent, privacy, and attention to skin integrity. The nurse should avoid massage in areas with wounds, infection, significant swelling, vascular complications, or other conditions in which massage may be inappropriate.

Hand Massage

Hand massage involves gentle manipulation of the hands and may provide sensory stimulation and relaxation. Because the hands are readily accessible, the technique can be relatively easy to administer in a hospital setting.

The nurse should assess for wounds, intravenous access, inflammation, swelling, or other conditions that may make massage inappropriate.

Mechanisms Proposed for Massage and Pain Relief

Massage may influence pain through several proposed mechanisms. Mechanical stimulation of the skin and underlying tissues provides non-painful sensory input that may influence transmission and perception of pain. Massage may also promote relaxation and reduce muscle tension.

The psychological effects of human touch may also contribute to comfort. Feeling cared for and supported may reduce anxiety and improve the mother's ability to cope with pain.

However, massage should be considered a complementary intervention rather than a substitute for appropriate medical evaluation or prescribed analgesia.

Nursing Role in Non-Pharmacological Pain Management

The nurse has a significant role in selecting, implementing, and evaluating non-pharmacological interventions.

The nursing process can be applied:

Assessment

Assess pain intensity, location, quality, duration, aggravating factors, maternal preference, skin integrity, mobility, and relevant contraindications.

Planning

Select an appropriate intervention based on the mother's condition and preference. Establish a realistic goal, such as improved comfort or reduced pain intensity.

Intervention

Explain the procedure, obtain consent, maintain privacy, perform hand hygiene, provide the selected intervention safely, and observe the mother's response.

Evaluation

Reassess pain after the intervention using the same pain assessment method where possible. Compare the post-intervention score with the baseline score and document the response.

Foot and Hand Massage as a Complementary Nursing Intervention

Foot and hand massage can be considered complementary comfort measures for post-caesarean mothers. These interventions are relatively simple and can be performed without sophisticated equipment when appropriately trained personnel are available.

Massage should be performed gently and systematically. The mother should be positioned comfortably. The nurse should explain the procedure and obtain consent before beginning.

A typical session may begin with gentle stroking to promote relaxation. Gradual pressure can then be applied according to the mother's comfort. The nurse should continuously monitor verbal and non-verbal responses and stop the intervention if discomfort occurs.

The duration and frequency of massage should be determined according to the study protocol or institutional practice. In research, intervention procedures should be standardized so that participants receive comparable treatment.

For a quasi-experimental study evaluating foot and hand massage, baseline pain intensity should be measured before the intervention. Following the intervention, pain intensity should be reassessed using the selected standardized pain scale.

Foot Massage and Pain Reduction

Foot massage is a complementary nursing intervention in which systematic manual stimulation is applied to the feet using gentle stroking, rubbing, kneading, and pressure according to the individual's comfort and the therapeutic purpose. In postoperative nursing care, foot massage may be used as an adjunct to prescribed analgesic treatment to promote relaxation, improve comfort, and support the patient's ability to cope with pain. For post-caesarean mothers, the intervention is particularly relevant because the mother may experience abdominal discomfort, restricted mobility, fatigue, anxiety, and difficulty performing routine activities during the early postoperative period. A simple intervention such as foot massage can provide an opportunity for comfort while also giving the nurse an active role in individualized pain management.

Post-caesarean pain is primarily associated with surgical tissue injury, but the experience of pain is influenced by several additional factors. Anxiety, fear, fatigue, lack of sleep, unfamiliar surroundings, and concern about caring for the newborn may intensify the subjective experience of pain. Foot massage may contribute to pain management by providing a pleasant sensory stimulus and promoting a state of physical and psychological relaxation. When the mother becomes more relaxed, muscle tension and anxiety may decrease, which may in turn influence the way pain is perceived and tolerated.

Foot massage does not directly repair the surgical wound or remove the underlying cause of postoperative pain. Rather, it should be understood as a complementary intervention that may modify the patient's perception of discomfort and improve overall comfort. It should therefore be integrated into a comprehensive pain-management plan rather than being used as a replacement for clinically indicated analgesic therapy. The nurse should assess the mother's pain before the intervention, identify contraindications, explain the procedure, obtain consent, perform the massage appropriately, and reassess pain after completion.

The procedure begins with appropriate preparation of the mother and the environment. The mother should be positioned comfortably, usually in a position that does not place unnecessary pressure on the abdominal incision. Privacy should be maintained, and the environment should be quiet and comfortable. Hand hygiene should be performed before and after the procedure. The nurse should inspect the feet for skin lesions, wounds, inflammation, infection, swelling, impaired circulation, or other conditions that may make massage inappropriate. The nurse should also consider any clinical restrictions or medical conditions identified by the healthcare team.

Gentle stroking is commonly used at the beginning of massage to familiarize the mother with the touch and promote relaxation. The nurse may gradually progress to gentle rubbing or kneading while maintaining communication with the mother regarding pressure and comfort. The pressure should remain appropriate and should never cause pain. The mother should be encouraged to communicate if the pressure is uncomfortable or if any unusual sensation occurs.

The therapeutic value of foot massage may also arise from the opportunity it provides for supportive nurse–patient interaction. During massage, the nurse can communicate with the mother, provide reassurance, answer questions, and reinforce positive coping strategies. This interaction may be particularly meaningful for a mother who feels vulnerable after surgery. Thus, foot massage can have both physical and psychological dimensions.

The response to foot massage should be evaluated objectively whenever possible. A standardized pain assessment tool, such as a numerical rating scale, may be used before and after the intervention. Other outcomes such as relaxation, comfort, willingness to move, ability to rest, and satisfaction with care may also be documented. In a research study, the same assessment tool and assessment procedure should be used consistently to minimize measurement variation.

The timing of foot massage is another consideration. The intervention may be provided during a period when the mother is experiencing pain or when routine nursing care permits uninterrupted relaxation. In a research setting, the timing, duration, technique, and frequency of massage should be standardized. Standardization helps ensure that differences in pain outcomes can be interpreted in relation to the intervention rather than differences in how the intervention was delivered.

Foot massage may also be particularly acceptable because it does not directly involve the abdominal surgical site. A mother may be reluctant to have the area around her incision touched because of tenderness or fear of disrupting the wound. Foot massage provides a distal approach to comfort while avoiding manipulation of the surgical wound.

An important nursing principle is that complementary interventions should be patient-centered. Not every mother will prefer massage. Some may feel uncomfortable with touch, while others may prefer relaxation exercises, music, positioning, or other measures. The nurse should respect the mother's

preference and cultural expectations. Consent should be obtained, and the intervention should be stopped if the mother reports discomfort.

Foot massage may therefore be viewed as a supportive component of holistic postoperative nursing care. Its purpose is not simply to reduce a numerical pain score but to promote comfort, relaxation, confidence, and participation in recovery. When combined with appropriate assessment, prescribed analgesics, mobility support, breastfeeding assistance, and psychological support, foot massage can form part of an individualized maternal pain-management strategy.

Hand Massage and Pain Reduction

Hand massage is another simple complementary intervention that can be incorporated into nursing care for patients experiencing acute pain. It involves gentle manipulation and stimulation of the hands using techniques such as stroking, rubbing, and gentle kneading. The hands are easily accessible, and the intervention can generally be performed without requiring complex equipment. For post-caesarean mothers, hand massage can provide a convenient method of promoting comfort and relaxation while avoiding direct manipulation of the abdominal surgical area.

The experience of postoperative pain involves both physiological and psychological processes. Surgical stimulation activates nociceptive pathways, while anxiety and emotional distress may increase the patient's perception of pain. Hand massage provides a form of non-painful sensory stimulation and may encourage the mother to focus on a pleasant physical sensation rather than continuously concentrating on postoperative discomfort.

The simplicity of hand massage is one of its practical advantages. A nurse who has received appropriate training can perform the intervention at the bedside. The mother does not necessarily need to change position significantly, which may be beneficial when movement is uncomfortable during the early postoperative period. The intervention can therefore be integrated into routine nursing care while maintaining appropriate safety precautions.

Before performing hand massage, the nurse should assess the condition of both hands. The presence of intravenous cannulas, wounds, swelling, inflammation, infection, skin breakdown, or other abnormalities should be considered. If an intravenous line is present, the nurse should avoid applying pressure that could displace or interfere with the device. Massage should also be avoided or modified when a clinical condition makes manual stimulation inappropriate.

The mother should be informed about the purpose and procedure. A clear explanation can reduce anxiety and improve cooperation. The nurse should ask about preferred pressure and should continuously observe the mother's response. Gentle pressure is generally more appropriate than vigorous manipulation in a postoperative patient.

The massage may begin with light stroking across the palm and back of the hand. The fingers can be gently manipulated one at a time, followed by gentle circular movements over appropriate areas of the hand. The wrist may also receive gentle movement when clinically appropriate. The technique should remain comfortable and should never produce pain, numbness, tingling, or other concerning symptoms.

Hand massage may have an important psychological dimension. Human touch, when welcomed by the patient, can communicate attention, reassurance, and emotional support. A post-caesarean mother may experience fear regarding her recovery, concerns about the newborn, and anxiety about postoperative activities. Supportive touch may help create a sense of comfort and security.

The nurse should recognize that hand massage is not intended to replace analgesic medication. If the mother reports moderate or severe postoperative pain, appropriate clinical assessment and prescribed treatment should not be delayed. Complementary interventions are best used alongside appropriate medical and nursing management.

Pain should be assessed before and after hand massage. A numerical rating scale or another validated pain assessment tool may be used. Documentation should include the pre-intervention

pain score, intervention provided, duration of intervention, patient's response, and post-intervention pain score. Consistent documentation is especially important in research studies.

Hand massage may also be combined with breathing exercises. The mother can be encouraged to breathe slowly and comfortably while the nurse provides gentle massage. This combination may promote relaxation and may make the intervention more meaningful to the patient.

The intervention can also be used during periods when the mother is resting. However, the nurse should avoid unnecessary disturbance if the mother is sleeping comfortably. Pain assessment and clinical needs should guide the timing of the intervention.

In a research context, hand massage should be delivered according to a clearly defined protocol. The protocol may specify the sequence of massage movements, duration, frequency, pressure, body position, and environmental conditions. Standardization is essential because variations in technique can influence the intervention's effects.

Hand massage may therefore represent a feasible and patient-centered complementary intervention in post-caesarean nursing. Its accessibility, low equipment requirement, and potential to promote relaxation make it an intervention of interest in nursing research.

Combined Foot and Hand Massage Intervention

Combined foot and hand massage refers to the systematic application of massage techniques to both the hands and feet during the same intervention period. The concept is based on providing sensory stimulation to two easily accessible peripheral body regions while simultaneously promoting relaxation and comfort. In the context of post-caesarean pain management, combined massage may provide a broader relaxation experience than intervention involving only one body region.

A post-caesarean mother often experiences pain together with fatigue, anxiety, reduced mobility, and sleep disturbance. Because the abdominal surgical area may be sensitive, massage can be provided to non-incisional areas such as the hands and feet. This approach may allow the nurse to provide therapeutic touch without directly manipulating the surgical wound.

The combined intervention should follow a standardized sequence. The mother should first be assessed for pain and general condition. The nurse should explain the procedure and obtain consent. The hands and feet should be inspected for contraindications. The mother should then be positioned comfortably, and the intervention can begin.

A standardized protocol is particularly important when combined foot and hand massage is used as part of a research study. For example, the protocol can specify whether hand massage is performed first followed by foot massage, or whether the reverse sequence is used. The duration of each component should also be predetermined. The pressure and techniques should be clearly described so that the intervention can be reproduced consistently.

The environment can influence relaxation. A quiet and comfortable environment may help the mother concentrate on the intervention. Unnecessary interruptions should be minimized where possible. Privacy should be maintained throughout the procedure.

The nurse may begin with gentle stroking to establish comfort. Gradually, gentle rubbing and kneading movements may be introduced. Communication should be maintained throughout the procedure, but excessive conversation should be avoided if the mother prefers quiet relaxation.

Combined massage may also provide an opportunity to teach the mother simple self-relaxation techniques. The nurse can encourage slow breathing and explain how relaxation can support coping with postoperative discomfort. Education should be brief and appropriate to the mother's level of fatigue.

The effectiveness of combined foot and hand massage should be evaluated using measurable outcomes. Pain intensity is the primary outcome in many studies. Additional outcomes may include anxiety, comfort, sleep, satisfaction, mobility, or need for rescue analgesics. However, each outcome should be measured using an appropriate and validated method.

One of the important principles of nursing research is the use of baseline measurements. Pain intensity should be assessed before the intervention. After the intervention, the pain should be reassessed using the same scale and under comparable conditions. This allows researchers to determine the change in pain intensity.

Combined massage may also have a role in patient education. Mothers can be informed that complementary approaches may support comfort but should not replace prescribed treatment. They should also be taught when to report increasing pain, wound problems, fever, excessive bleeding, or other concerning symptoms.

Safety is an important component of the intervention. Massage should not be performed over areas of infection, open wounds, severe inflammation, or other contraindicated conditions. The nurse should also be alert to conditions associated with impaired circulation or other risks where massage may not be appropriate.

The combined intervention is especially relevant to nursing because it can potentially be implemented without expensive equipment. However, feasibility should not be confused with universal appropriateness. Clinical assessment remains necessary before the intervention is provided.

From a research perspective, combined foot and hand massage can be evaluated using a quasi-experimental design. Mothers may be allocated to experimental and comparison groups according to the study design. The experimental group receives the standardized massage intervention in addition to routine care, while the comparison group receives routine care or another specified intervention. Pain scores before and after intervention can then be compared.

Mechanisms of Pain Relief through Massage

The mechanisms through which massage may influence pain are complex and involve physiological, neurological, psychological, and behavioral processes. Massage should not be viewed as acting through a single mechanism. Rather, several mechanisms may operate simultaneously.

One proposed mechanism involves sensory stimulation. Gentle pressure and movement stimulate sensory receptors in the skin and underlying tissues. Non-painful sensory input may influence the processing of nociceptive signals within the nervous system. This concept is consistent with the broader gate-control model of pain, which proposes that activity in non-nociceptive sensory pathways can modulate the transmission of pain-related signals at the spinal level.

The gate-control concept has been influential in explaining why non-painful stimulation, such as rubbing an injured area, can sometimes temporarily alter pain perception. Massage provides continuous sensory stimulation and may therefore contribute to modulation of the pain experience. However, pain physiology is complex, and the gate-control theory alone does not fully explain all effects of massage.

Another potential mechanism is reduction in muscle tension. Pain and anxiety can produce protective muscle contraction. Increased muscle tension may contribute to discomfort and can make movement more difficult. Gentle massage may promote relaxation and reduce perceived muscular tension. When relaxation improves, the individual may feel more comfortable and may become more willing to move.

Massage may also influence psychological responses to pain. Anxiety and fear can amplify the subjective experience of pain. A calm environment, supportive touch, and focused attention may reduce anxiety and improve coping. For a post-caesarean mother, emotional reassurance can be particularly important because she is simultaneously recovering from surgery and adapting to newborn care.

The relaxation response may also influence autonomic activity. Gentle massage may be associated with reduced physiological arousal in some individuals. The patient may experience slower

breathing, a sense of calm, and improved subjective comfort. These responses can indirectly influence pain tolerance.

Another proposed mechanism is distraction. During massage, the patient's attention may be directed toward the sensory experience of touch. This may reduce the amount of attention devoted to pain. Attention is an important component of pain perception because pain is not experienced independently of cognitive processes.

Massage may also contribute to a sense of control. Postoperative patients may feel that pain and recovery are outside their control. Participating in a structured comfort intervention can provide an opportunity for active involvement in recovery. A mother may feel more confident when she learns that several methods are available to help her cope with discomfort.

The therapeutic relationship may also contribute to the effect. Nursing care is not solely technical; interpersonal interaction is an important component of holistic care. A respectful and supportive interaction may decrease distress and increase satisfaction with care.

The biochemical mechanisms proposed in some massage literature include changes in stress-related physiological responses and neurochemical pathways. However, these mechanisms should be described cautiously because evidence varies depending on the population, massage technique, outcome measured, and research design.

Massage should therefore be understood as a multidimensional intervention. Its potential effect on pain may arise from sensory stimulation, relaxation, reduction in muscle tension, distraction, psychological support, and changes in the patient's perception of control.

Evidence from Previous Studies

Research into massage therapy has examined its potential role in managing pain and anxiety in various populations, including surgical patients, women during childbirth, postpartum women, and patients experiencing different types of acute or chronic pain. The findings across studies suggest that massage may provide short-term improvements in subjective comfort and pain for some patients, although the magnitude and consistency of effects vary.

Studies examining postoperative patients have frequently focused on non-pharmacological interventions as adjuncts to standard analgesic treatment. Massage has been investigated because it is relatively inexpensive, does not require complex equipment, and can be integrated into nursing care. In many studies, researchers have measured pain before and after the intervention using numerical or visual pain scales.

Research involving foot massage has reported reductions in self-reported pain in some clinical populations. Proposed explanations include sensory stimulation, relaxation, distraction, and reduction of anxiety. However, study findings should be interpreted according to methodological characteristics such as sample size, intervention duration, control condition, and pain measurement method.

Studies of hand massage have similarly explored its effects on pain and anxiety. Hand massage is particularly suitable for bedside nursing because the hands are accessible and the intervention can be delivered without requiring extensive repositioning. Some studies have reported improvements in comfort and relaxation following hand massage, although variations in protocols make direct comparison between studies difficult.

Research among postoperative patients has also investigated massage as an adjunct rather than an alternative to medication. This distinction is important. Ethical and clinically appropriate research should generally ensure that participants receive necessary routine care while the complementary intervention is evaluated.

In maternal health research, non-pharmacological pain-management approaches have received attention because postpartum women may experience pain while simultaneously caring for their

newborns. Interventions that promote relaxation and comfort may support maternal functioning. Massage has therefore been explored in relation to pain, anxiety, satisfaction, and overall comfort.

The evidence should nevertheless be interpreted carefully. A reduction in pain score immediately after massage does not necessarily demonstrate a long-term effect on recovery. Pain is influenced by multiple variables, including medication administration, time since surgery, physical activity, sleep, anxiety, and individual differences.

Another important consideration is the placebo or attention effect. Patients receiving an intervention may receive additional attention from healthcare personnel, which itself can influence subjective outcomes. In a controlled research design, researchers should therefore attempt to distinguish the specific effect of massage from the general effect of attention and supportive care.

The methodological quality of individual studies is also important. Randomized controlled trials generally provide stronger evidence for causal inference than uncontrolled studies. Quasi-experimental studies can provide useful evidence in clinical settings, particularly when random allocation is impractical, but potential differences between groups must be considered.

Sample size is another factor affecting interpretation. Small studies may demonstrate promising results but may have limited statistical power and generalizability. Larger studies with clearly described intervention protocols can provide more reliable estimates of effect.

Pain measurement also influences findings. Subjective pain scales are appropriate because pain is a subjective experience, but scores may vary according to mood, expectations, time of day, medication, and environmental conditions. Therefore, studies should use validated instruments and standardized assessment procedures.

Research protocols should clearly describe the massage intervention. Important details include the body area treated, massage technique, duration, frequency, pressure, timing relative to surgery, provider training, and comparison intervention. Without such details, replication and comparison between studies become difficult.

For a quasi-experimental study among post-caesarean mothers, evidence from previous research provides a rationale for investigating whether a standardized foot and hand massage intervention is associated with changes in pain intensity. The study can contribute additional evidence specific to the selected maternal population and clinical setting.

Research Gap

Although complementary therapies such as massage have been examined in different patient populations, findings may not automatically be generalized to post-caesarean mothers in every clinical setting. Differences in cultural expectations, healthcare practices, analgesic protocols, maternal characteristics, and intervention procedures may influence outcomes.

A study conducted among post-caesarean mothers in a selected maternity hospital in Noida, Uttar Pradesh, can therefore provide setting-specific evidence. The study can assess whether the intervention is feasible and whether measurable changes in pain intensity occur under the conditions specified in the research protocol.

The study can also contribute to nursing practice by examining an intervention that may be delivered by appropriately trained nursing personnel as a complement to routine postoperative care.

Quasi-Experimental Approach to Evaluating Massage

A quasi-experimental approach is an important research design for evaluating the effectiveness of nursing interventions when random allocation of participants is not feasible or practical. In maternal and newborn nursing, quasi-experimental designs can be used to examine whether a planned nursing intervention produces measurable changes in patient outcomes under real clinical conditions. In the context of post-caesarean pain management, a quasi-experimental study can be designed to evaluate whether foot and hand massage is associated with a reduction in pain intensity

among mothers following caesarean birth. Such an approach provides an opportunity to evaluate a complementary nursing intervention while mothers continue to receive routine postoperative care.

The proposed study, **"A Quasi-Experimental Study to Assess the Effectiveness of Foot and Hand Massage in Reduction of Pain Intensity among Post-Caesarean Mothers in a Selected Maternity Hospital at Noida, Uttar Pradesh,"** focuses on pain intensity as the principal outcome. Postoperative pain is a subjective experience and can be influenced by several factors, including the type of surgery, time since surgery, analgesic administration, anxiety, previous experiences, fatigue, sleep, mobility, and individual pain perception. Therefore, a research design that permits systematic measurement of pain before and after an intervention is useful for determining whether a clinically meaningful change occurs.

In a quasi-experimental study, participants are generally assigned to an experimental group and a comparison or control group without true randomization. The experimental group receives the intervention under investigation, while the control group receives routine care or another specified condition. The two groups can then be compared in relation to changes in the selected outcome. When studying massage for post-caesarean pain, the experimental group may receive standardized foot and hand massage in addition to routine postoperative care, whereas the comparison group may receive routine postoperative care according to the hospital's usual practice.

The use of a comparison group is important because postoperative pain naturally changes over time. A mother may experience a reduction in pain several hours after surgery even without massage because of tissue recovery, medication, rest, or gradual adaptation. If pain is measured only before and after massage in a single group, it may be difficult to determine how much of the change is specifically associated with the intervention. A comparison group provides a reference against which the observed change can be examined.

A pre-test and post-test approach can be particularly useful for this type of research. In this approach, pain intensity is measured before the intervention and again after the intervention. The difference between the pre-intervention and post-intervention scores can be calculated. Changes within the experimental group can then be compared with changes in the comparison group. This approach allows the researcher to determine whether the reduction in pain is greater in the intervention group than would be expected from routine postoperative recovery alone.

The selection of participants is another important component of the quasi-experimental approach. The researcher should establish clearly defined inclusion and exclusion criteria before recruitment begins. Inclusion criteria may specify factors such as post-caesarean status, age range, postoperative period, ability to communicate, and willingness to participate. Exclusion criteria may include conditions in which massage is clinically inappropriate or circumstances that could interfere with accurate pain assessment.

The timing of intervention is particularly important in postoperative research. Pain intensity can change substantially during the hours following surgery. Therefore, the researcher should identify a consistent postoperative period for baseline assessment and intervention. For example, all participants may be assessed within a predetermined postoperative period according to the study protocol. Standardization of timing reduces variation and improves comparability between participants.

The intervention itself should be clearly operationalized. A research protocol should specify exactly what is meant by foot and hand massage. It may describe the sequence of hand and foot massage, type of strokes, pressure, duration, frequency, positioning, and environmental conditions. The person delivering the intervention should receive appropriate training to ensure that the technique is applied consistently.

Standardization is essential because massage is not a single uniform intervention. Different techniques and durations may produce different responses. If one participant receives gentle stroking for a short period and another receives vigorous kneading for a longer period, the

intervention is not sufficiently standardized for rigorous comparison. A written intervention protocol helps reduce this variation.

The researcher should also determine how pain intensity will be measured. A validated pain assessment tool should be selected and used consistently. A Numerical Rating Scale, for example, can allow participants to report their pain using a defined numerical range. The same scale should be used for both pre-intervention and post-intervention assessments whenever possible.

Pain assessment should be conducted in a manner that minimizes unnecessary influence from the researcher. Participants should be encouraged to report the pain they actually experience rather than what they believe the researcher expects. Clear instructions should be provided, and the meaning of the scale should be explained before the first assessment.

The reliability and validity of research instruments are important considerations. If the researcher uses a pain assessment tool, the tool should be appropriate for the study population. If additional instruments are used to measure demographic characteristics, anxiety, satisfaction, or other outcomes, their reliability and validity should also be established or supported by appropriate evidence.

Ethical considerations are fundamental to quasi-experimental research involving post-caesarean mothers. Ethical approval should be obtained from the appropriate institutional review or ethics committee before data collection begins. Participants should be informed about the purpose of the study, procedures involved, expected benefits, potential discomforts, confidentiality, voluntary participation, and their right to withdraw according to the approved protocol.

Informed consent is particularly important when the intervention involves physical touch. The mother should understand that massage is voluntary and that declining participation will not affect her routine care. The researcher should respect privacy, dignity, cultural preferences, and individual boundaries.

Confidentiality should be maintained throughout the research process. Participant names should preferably be replaced by identification codes in research records. Data should be stored securely and should be accessible only to authorized persons. Research findings should be presented in aggregate form so that individual participants cannot ordinarily be identified.

The researcher must also ensure that the intervention does not interfere with necessary postoperative care. Foot and hand massage should be considered complementary to routine care rather than a substitute for prescribed analgesics or medical treatment. If a participant experiences severe or unexpected pain, the appropriate clinical assessment and management should take priority over research procedures.

The study should also have a clearly defined procedure for handling adverse events. Although gentle massage may be a relatively simple intervention, the researcher should monitor for discomfort, skin irritation, dizziness, worsening pain, or any other unexpected response. The intervention should be stopped if clinically indicated, and appropriate medical or nursing care should be provided.

Data collection should be systematic. Demographic and clinical variables that may influence pain should be recorded according to the study protocol. These may include age, parity, type of caesarean section, previous caesarean history, postoperative duration, analgesic administration, and other relevant characteristics. Recording these variables helps the researcher understand the characteristics of the study population.

The statistical analysis should correspond to the research questions and design. Descriptive statistics can be used to summarize demographic and clinical characteristics. Measures such as frequency, percentage, mean, and standard deviation may be used depending on the nature and distribution of the data. Inferential statistical tests can then be selected according to the research design, measurement scale, distribution, and assumptions of the data.

Within-group comparisons can examine whether pain scores changed from pre-intervention to post-intervention. Between-group comparisons can examine whether the amount of change differed between the experimental and comparison groups. The statistical method should be selected in consultation with an appropriate statistical expert and should be specified before analysis whenever possible.

The researcher should distinguish statistical significance from clinical significance. A statistically significant difference does not necessarily mean that the change is large enough to be meaningful to the mother. Conversely, a clinically meaningful improvement may not reach statistical significance in a small sample. Therefore, interpretation should consider both the magnitude of change and the statistical evidence.

A quasi-experimental study can provide valuable evidence, but it also has limitations. Because participants are not randomly allocated, the two groups may differ in important characteristics before the intervention. Selection bias may therefore influence the findings. The researcher can reduce this concern by using clearly defined eligibility criteria, obtaining baseline measurements, examining group characteristics, and applying appropriate statistical methods.

Another limitation is the possibility of confounding factors. Analgesic medication, mobility, time since surgery, breastfeeding, sleep, and emotional state may influence pain independently of massage. Researchers should record relevant variables and maintain consistent procedures whenever feasible.

Researcher bias is another consideration. If the person delivering the intervention also assesses the outcome, expectations may influence assessment. Blinding may be difficult in massage studies because participants generally know whether they received massage. Nevertheless, standardized procedures and objective documentation can reduce potential bias.

Despite these limitations, a well-designed quasi-experimental study can provide useful clinical evidence. It can demonstrate whether a standardized massage intervention is feasible in a maternity setting and whether measurable changes in pain intensity occur under specified clinical conditions.

Clinical Application among Post-Caesarean Mothers

The clinical application of foot and hand massage begins with recognition that post-caesarean pain management should be individualized. Every mother does not experience pain in the same manner, and the severity and characteristics of postoperative discomfort can vary. The nurse should therefore assess the mother before selecting a complementary intervention.

The first step is a comprehensive postoperative assessment. The nurse should assess vital signs, level of consciousness, surgical wound, bleeding, mobility, pain intensity, prescribed medication, and general condition. The mother should also be asked about the location and quality of pain and factors that increase or decrease discomfort.

The mother should be informed about the purpose of massage. The nurse can explain that foot and hand massage is intended to promote relaxation and comfort and may complement prescribed pain treatment. The nurse should avoid making unrealistic promises regarding pain reduction.

Consent should be obtained before touching the mother. The mother should be allowed to decline or stop the intervention at any time. Respect for autonomy is particularly important during postpartum care because mothers may already feel physically vulnerable following surgery.

Before foot massage, the nurse should inspect the feet. Areas of skin breakdown, infection, significant swelling, wounds, or other abnormalities should be identified. The nurse should consider the mother's overall clinical condition and follow institutional guidance regarding contraindications.

Similarly, beforehand massage, the nurse should inspect the hands. Particular attention should be given to intravenous access sites, wounds, swelling, inflammation, or other conditions. Massage should be modified or avoided where clinically indicated.

The mother should be positioned comfortably. Positioning should avoid pressure on the abdominal incision and should permit the nurse to access the hands and feet without causing unnecessary movement. Pillows or other appropriate supports may be used.

The environment should be conducive to relaxation. Excessive noise and unnecessary interruptions should be minimized where possible. Adequate privacy should be maintained. The mother should feel safe and comfortable during the procedure.

The massage should begin gently. Initial light stroking allows the mother to become accustomed to the intervention. The nurse should observe the mother's response and ask whether the pressure is comfortable.

For foot massage, gentle movements can be applied systematically to appropriate areas of the foot. The technique should remain comfortable and should not produce pain. Excessive pressure should be avoided, especially in a postoperative patient.

For hand massage, gentle stroking and rubbing can be applied to the palms, backs of the hands, and fingers according to the standardized protocol. If an intravenous device is present, the nurse should ensure that it is protected and not disturbed.

The sequence and duration of the intervention should be standardized when massage is being used as part of a research study. This may include a predetermined duration for hand massage followed by foot massage or another sequence specified in the research protocol.

The nurse should continuously monitor the mother's verbal and non-verbal response. If the mother reports increased pain, dizziness, discomfort, numbness, or any unusual symptom, the procedure should be stopped and the mother should be assessed.

The nurse can encourage slow, comfortable breathing during the massage. Breathing should not be forced. The combination of gentle massage and relaxed breathing may help the mother focus on comfort and reduce tension.

After completion of massage, the mother's pain should be reassessed using the same standardized tool used before the intervention. The nurse should also ask whether she feels more relaxed or comfortable.

Documentation is essential. The nurse should record the intervention, duration, patient's response, pain score before and after the intervention, and any relevant observations.

Foot and hand massage should be integrated into the overall plan of care. If the mother requires analgesic medication, the prescribed medication should be administered according to clinical orders. Massage should not delay treatment of significant pain.

Clinical application also requires consideration of breastfeeding. Post-caesarean mothers may experience difficulty positioning themselves comfortably during breastfeeding. Pain management should therefore be coordinated with breastfeeding assistance. The nurse can help the mother find a position that minimizes abdominal pressure.

Mobility is another important consideration. Adequate pain management may make it easier for the mother to participate in gradual movement. The nurse should follow postoperative mobility recommendations and assist the mother appropriately.

The clinical goal is not simply to achieve a lower pain score. Effective pain management should support functional recovery. A mother who can turn in bed, sit comfortably, stand, walk safely, breastfeed, hold her baby, and rest adequately may be experiencing meaningful improvement even when some pain remains.

Implications for Maternal and Newborn Nursing

The use of complementary pain-management interventions has several implications for maternal nursing practice. Nurses are frequently the healthcare professionals who spend the greatest amount

of time with postpartum mothers and therefore have an important role in identifying pain and implementing appropriate comfort measures.

Pain assessment should be incorporated into routine maternal nursing care. Pain should not be considered an unavoidable consequence of caesarean birth that requires no intervention. Instead, nurses should assess pain systematically and respond according to the mother's needs.

A holistic approach to pain management recognizes that physical pain is interconnected with emotional and social experiences. A mother may be worried about her newborn, concerned about her recovery, or anxious about breastfeeding. These concerns can influence her perception of pain. Nursing care should therefore address both physical and psychological needs.

Therapeutic massage may provide an additional nursing intervention that promotes patient comfort. Because it can be performed at the bedside with minimal equipment, it may be feasible in a range of maternity settings when appropriate training and clinical safeguards are available.

Nursing education should include knowledge of non-pharmacological pain-management approaches. Student nurses and practicing nurses should understand the principles, benefits, limitations, contraindications, and safety precautions associated with therapeutic massage.

Nursing educators can incorporate massage and other complementary interventions into simulation-based learning. Students can practice communication, consent, positioning, technique, assessment, and documentation before applying the intervention in clinical settings.

The intervention also has implications for patient education. Mothers can be taught that pain management may involve several approaches. They can be encouraged to report pain early rather than waiting until it becomes severe.

Patient education can also help mothers understand that analgesic medication and complementary methods are not necessarily competing approaches. Appropriate non-pharmacological methods may complement prescribed treatment.

The intervention may have implications for nursing administration. If a maternity unit decides to incorporate therapeutic massage into routine care, administrators may need to develop protocols addressing staff training, patient eligibility, contraindications, documentation, infection prevention, and quality monitoring.

Institutional protocols can improve consistency. A standardized protocol may specify who can provide massage, how the intervention should be performed, when it should be avoided, and how patient response should be documented.

Nursing administration should also ensure that complementary interventions do not replace evidence-based medical treatment. The availability of massage should not result in delayed analgesia or inadequate assessment of severe postoperative pain.

Maternal and newborn nursing is fundamentally family-centered. Improving maternal comfort can potentially support the mother's participation in newborn care. A mother who is more comfortable may find it easier to hold, feed, communicate with, and care for her infant.

However, nurses should avoid assuming that pain reduction automatically results in improved maternal–infant outcomes. These relationships should be evaluated through appropriate research rather than presumed.

The intervention also has implications for patient satisfaction. Respectful communication, attention, comfort measures, and individualized care can influence the mother's perception of her hospital experience. Massage may provide one opportunity for personalized nursing interaction.

Culturally sensitive care is essential. Attitudes toward physical touch vary among individuals and communities. Nurses should ask permission and respect preferences rather than assuming that massage is universally acceptable.

The economic implications may also be relevant. Foot and hand massage generally does not require expensive equipment. If shown to be effective and feasible, it may represent a relatively low-resource complementary intervention. Nevertheless, training and staff time have associated costs and should be considered when evaluating implementation.

Research implications are also significant. Nursing researchers can investigate massage in relation to pain intensity, anxiety, sleep quality, maternal satisfaction, mobility, breastfeeding experience, and analgesic requirements. Such research can expand understanding of how complementary interventions may fit within comprehensive postoperative care.

Future research should use clearly described intervention protocols. Studies should report massage duration, frequency, technique, provider training, timing after surgery, comparison condition, pain measurement instrument, and relevant clinical variables.

Researchers should also consider larger sample sizes and appropriate study designs. Randomized controlled trials may provide stronger causal evidence where feasible, while well-designed quasi-experimental studies can provide useful evidence in real-world clinical environments.

Conclusion and Future Directions

Post-caesarean pain remains an important issue in maternal nursing because it can affect physical recovery, mobility, breastfeeding, sleep, psychological well-being, and participation in newborn care. Effective pain management requires systematic assessment and a combination of appropriate pharmacological and non-pharmacological approaches.

Foot and hand massage represent complementary interventions that may support comfort and relaxation among post-caesarean mothers. Their potential usefulness is based on several possible mechanisms, including sensory stimulation, relaxation, reduction of muscle tension, distraction, emotional support, and improved coping with pain.

A quasi-experimental approach provides a practical method for evaluating the effectiveness of such an intervention in a clinical maternity setting. By measuring pain before and after intervention and comparing outcomes with a suitable comparison group, researchers can obtain information about whether the intervention is associated with a measurable reduction in pain intensity.

A strong research protocol is essential. Participant selection, timing of intervention, massage technique, duration, pain measurement, routine care, analgesic administration, and data collection should be clearly defined. Consistency in implementation strengthens the credibility of the findings.

Clinical application requires careful patient assessment and attention to safety. Massage should not be performed automatically for every post-caesarean mother. The nurse should assess the patient's clinical condition, identify contraindications, obtain consent, maintain infection-prevention practices, and monitor the response throughout the procedure.

The intervention should remain complementary to standard postoperative management. Severe or unexpected pain requires appropriate clinical assessment and treatment. Massage should never delay necessary medical intervention or prescribed analgesia.

Future nursing research can examine whether repeated massage sessions produce sustained improvements compared with a single session. Researchers can also examine whether the intervention influences functional outcomes such as mobility, sleep, breastfeeding comfort, maternal satisfaction, and participation in newborn care.

Future studies could compare foot massage alone, hand massage alone, and combined foot and hand massage. Such comparisons could help determine whether combining the two interventions provides additional benefit compared with a single-site intervention.

Research could also examine different massage durations and frequencies. Determining the most practical duration for clinical implementation may be important because nursing staff have limited time and maternity units may have high patient workloads.

Further research should examine the influence of analgesic medication. Since postoperative mothers commonly receive pharmacological pain management, studies should document analgesic administration carefully when evaluating massage. This would help distinguish the effects associated with massage from changes associated with medication.

Larger multicenter studies could improve the generalizability of findings. A study conducted in one maternity hospital may provide valuable local evidence, but findings may not necessarily apply to all hospitals, populations, or healthcare systems. Multicenter research can provide evidence across different clinical environments.

Future studies may also incorporate patient-reported outcomes beyond pain intensity. A numerical pain score is important, but recovery involves multiple dimensions. Maternal comfort, anxiety, sleep, confidence, satisfaction, mobility, and ability to care for the newborn may provide a more comprehensive understanding of the intervention.

The development of standardized nursing protocols is another potential direction. If evidence demonstrates clinical usefulness, maternity hospitals may consider developing protocols for complementary massage interventions. Such protocols should include eligibility criteria, contraindications, informed consent, technique, duration, documentation, and evaluation.

Nursing education may also evolve to include greater emphasis on evidence-based complementary therapies. Undergraduate and postgraduate nursing programs can teach students how to critically evaluate complementary interventions and apply them safely within professional scope and institutional policy.

Technology may provide additional opportunities for future research. Digital pain diaries or electronic nursing records can facilitate repeated pain assessment and more detailed documentation of intervention response. Such systems may improve continuity of care and support research data collection.

Future research should also emphasize methodological quality. Researchers should use appropriate sample-size calculations, clearly defined outcomes, validated assessment instruments, standardized intervention procedures, transparent reporting, and suitable statistical methods.

The concept of patient-centered care should remain central. Massage should be offered as an option rather than imposed as a routine intervention. The mother's preferences, comfort, cultural expectations, clinical condition, and right to refuse should always be respected.

In maternal nursing, the ultimate purpose of pain management is not simply to eliminate every sensation of pain. The broader objective is to help the mother recover safely and comfortably so that she can participate in essential activities, interact with her newborn, establish breastfeeding when desired, rest adequately, and regain independence.

Foot and hand massage may contribute to this broader goal as part of a multimodal approach. Its value should be determined through systematic research rather than assumption. Evidence generated from well-designed quasi-experimental and controlled studies can help nurses understand when, how, and for whom therapeutic massage may be useful.

For the proposed study among post-caesarean mothers in a selected maternity hospital in Noida, Uttar Pradesh, the evaluation of foot and hand massage can provide clinically relevant information regarding its effect on pain intensity. The findings may contribute to the growing body of nursing evidence concerning complementary approaches to postoperative maternal care.

The future of maternal nursing increasingly emphasizes individualized, evidence-informed, compassionate, and holistic care. Within this framework, therapeutic massage represents one possible complementary intervention. When appropriately assessed, safely administered, systematically evaluated, and integrated with standard postoperative management, it may offer an additional avenue for promoting maternal comfort and supporting recovery after caesarean birth.

Ultimately, the successful management of post-caesarean pain requires collaboration among the mother, nurses, physicians, midwives, anesthetic team, lactation professionals, and family when appropriate. The nurse remains a key member of this team because of the nurse's continuous contact with the mother and central role in assessment, comfort measures, education, monitoring, and evaluation.

A comprehensive approach that combines appropriate analgesia, therapeutic positioning, mobility support, breastfeeding assistance, emotional support, relaxation techniques, and selected complementary interventions can provide a more complete model of postpartum care. Foot and hand massage can be considered within this broader framework rather than as an isolated treatment.

The continued development of high-quality nursing research will help clarify the role of therapeutic massage in post-caesarean recovery. Future evidence can guide clinical protocols, nursing education, patient education, and healthcare policy while ensuring that interventions remain safe, acceptable, and responsive to the needs of mothers and their newborns.

References

1. American College of Obstetricians and Gynecologists. (2024). *Optimizing postpartum care*. https://www.acog.org/clinical/clinical-guidance/committee-opinion/articles/2018/05/optimizing-postpartum-care
2. Apfelbaum, J. L., Chen, C., Mehta, S. S., & Gan, T. J. (2003). Postoperative pain experience: Results from a national survey suggest postoperative pain continues to be undermanaged. *Anesthesia& Analgesia, 97*(2), 534–540.
3. Chou, R., Gordon, D. B., de Leon-Casasola, O. A., Rosenberg, J. M., Bickler, S., Brennan, T., Carter, T., Cassidy, C. L., Chittenden, E. H., Degenhardt, E., Griffith, S., Manworren, R., McCarberg, B., Montgomery, R., Murphy, J., Perkal, M. F., Suresh, S., Sluka, K., Strassels, S., & Wu, C. L. (2016). Management of postoperative pain: A clinical practice guideline from the American Pain Society, the American Society of Regional Anesthesia and Pain Medicine, and the American Society of Anesthesiologists. *The Journal of Pain, 17*(2), 131–157. https://doi.org/10.1016/j.jpain.2015.12.008
4. Deussen, A. R., Ashwood, P., Martis, R., Stewart, F., & Grzeskowiak, L. E. (2023). Relief of pain due to uterine cramping/involution after birth. *Cochrane Database of Systematic Reviews*.
5. Hodnett, E. D., Gates, S., Hofmeyr, G. J., & Sakala, C. (2013). Continuous support for women during childbirth. *Cochrane Database of Systematic Reviews*, (7), CD003766. https://doi.org/10.1002/14651858.CD003766.pub5
6. International Association for the Study of Pain. (2020). *IASP announces revised definition of pain*. https://www.iasp-pain.org
7. Kwekkeboom, K. L., &Gretarsdottir, E. (2006). Systematic review of relaxation interventions for pain. *Journal of Nursing Scholarship, 38*(3), 269–277. https://doi.org/10.1111/j.1547-5069.2006.00113.x
8. McCaffery, M., &Pasero, C. (1999). *Pain: Clinical manual* (2nd ed.). Mosby.
9. Melzack, R., & Wall, P. D. (1965). Pain mechanisms: A new theory. *Science, 150*(3699), 971–979. https://doi.org/10.1126/science.150.3699.971
10. Moyer, C. A., Rounds, J., & Hannum, J. W. (2004). A meta-analysis of massage therapy research. *Psychological Bulletin, 130*(1), 3–18. https://doi.org/10.1037/0033-2909.130.1.3
11. National Institute for Health and Care Excellence. (2021). *Caesarean birth (NICE Guideline NG192)*. NICE. https://www.nice.org.uk/guidance/ng192
12. Pasero, C., & McCaffery, M. (2011). *Pain assessment and pharmacologic management*. Mosby/Elsevier.

13. Practice Guidelines for Obstetric Anesthesia: An Updated Report by the American Society of Anesthesiologists Task Force on Obstetric Anesthesia and the Society for Obstetric Anesthesia and Perinatology. (2016). *Anesthesiology, 124*(2), 270–300. https://doi.org/10.1097/ALN.0000000000000935
14. Smith, C. A., Levett, K. M., Collins, C. T., Dahlen, H. G., Ee, C. C., & Suganuma, M. (2018). Massage, reflexology and other manual methods for pain management in labour. *Cochrane Database of Systematic Reviews*, (3), CD009290. https://doi.org/10.1002/14651858.CD009290.pub3
15. Stevens, B. J., Abbott, L. K., Yamada, J., Harrison, D., Stinson, J., Taddio, A., Barwick, M., Latimer, M., Scott, S. D., Rashotte, J., Campbell-Yeo, M., & CIHR Team in Children's Pain. (2011). Epidemiology and management of painful procedures in children in Canadian hospitals. *Canadian Medical Association Journal, 183*(7), E403–E410. https://doi.org/10.1503/cmaj.101341
16. World Health Organization. (2022). *WHO recommendations for care of the preterm or low-birth-weight infant*. World Health Organization.
17. World Health Organization. (2022). *WHO recommendations on maternal and newborn care for a positive postnatal experience*. World Health Organization. https://www.who.int/publications/i/item/9789240045989